

Appendix F: Indicator Fact Sheet on HM depositions

(copy from the HELCOM web pages:

http://www.helcom.fi/environment2/ifs/ifs2006/en_GB/hmdepositions/)

Atmospheric depositions of heavy metals on the Baltic Sea

Editor(s): Alexey Gusev, EMEP MSC-E

1.1.1 Key message

😊 Total annual atmospheric depositions of heavy metals to the Baltic Sea have decreased in period from 1990 to 2004 by 51% for cadmium, 44% for mercury, and 69% for lead.

1.1.2 Results and Assessment

1.1.2.1 Relevance of the indicator for describing the developments in the environment

This indicator shows the levels and trends in cadmium, mercury, and lead atmospheric depositions to the Baltic Sea. The depositions of heavy metals represent the pressure of emission sources on the Baltic Sea aquatic environment.

1.1.2.2 Policy relevance and policy reference

HELCOM adopted a Recommendation in May 2001 for the cessation of hazardous substance discharges/emissions by 2020, with the ultimate aim of achieving concentrations in the environment near to background values for naturally occurring substances and close to zero for man-made synthetic substances.

1.1.2.3 Assessment

Total annual atmospheric depositions of heavy metals to the surface of the Baltic Sea have substantially decreased in period 1990-2004 (Figure 1). The most significant drop in depositions over the Baltic Sea is obtained for lead (69%). The decrease of cadmium and mercury depositions is amounted to 51% and 44%, respectively. On the level of individual sub-basins the most significant drop in cadmium depositions can be noted for the Gulf of Finland (68%). In case of lead the most significant decrease can be noted for the Gulf of Bothnia and the Gulf of Finland (73%). Largest decrease in mercury depositions is obtained for the Belt Sea (60%). In spatial distribution of heavy metals depositions on the Baltic Sea the highest levels can be noted for the southern-western part of the Baltic Sea (the Belt Sea and the Baltic Proper). Significant levels of lead and cadmium depositions can also be noted for the Gulf of Riga. Among the HELCOM countries the most significant contributions to depositions over the Baltic Sea belong to Poland, Germany, and Russia (Table 4).

The reduction of atmospheric input of lead, cadmium, and mercury to the Baltic Sea is a result of abatement measures as well as of economic contraction and industrial restructuring in Poland, Estonia, Latvia, Lithuania, and Russia in early 1990s.

1.1.2.4 Supporting information

http://www.msceast.org/HELCOM/HM_depositions_to_Baltic_Sea.pdf




Figure 1: Computed total annual atmospheric depositions of cadmium, mercury, and lead to the Baltic Sea for the period 1990-2004, (% of 1990).




Figure 2: Time-series of computed total annual atmospheric deposition of cadmium to six sub-basins of the Baltic Sea for the period 1990-2004 in tonnes/year as bars (left axis) and total deposition fluxes in $\text{g}/\text{km}^2/\text{year}$ as lines (right axis). Note that different scales are used for total depositions in tonnes/year and the same scales for total deposition fluxes. **Click image to enlarge!**




Figure 3: Time-series of computed total annual atmospheric deposition of lead to six sub-basins of the Baltic Sea for the period 1990-2004 in tonnes/year as bars (left axis) and total deposition fluxes in kg/km²/year as lines (right axis). Note that different scales are used for total depositions in tonnes/year and the same scales for total deposition fluxes. **Click image to enlarge!**




Figure 4: Time-series of computed total annual atmospheric deposition of mercury to six sub-basins of the Baltic Sea for the period 1990-2004 in tonnes/year as bars (left axis) and total deposition fluxes in $\text{g}/\text{km}^2/\text{year}$ as lines (right axis). Note that different scales are used for total depositions in tonnes/year and the same scales for total deposition fluxes. **Click image to enlarge!**

1.1.3 Data**Table 1.** Computed total annual depositions of cadmium to six Baltic Sea sub-basins for period 1990-2004. Units: tonnes/year

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004
GUB 2	1.4	1.4	1.4	1	1	1.3	0.8	1.1	1.1	1.3	1.1	0.9	0.9	0.8	
GUF 1.2	1	0.9	0.7	0.8	0.7	0.7	0.5	0.5	0.5	0.5	0.5	0.5	0.4	0.4	0.4
GUR 0.6	0.5	0.5	0.5	0.4	0.4	0.6	0.4	0.3	0.4	0.4	0.4	0.3	0.3	0.3	0.2
BAP 6.4	5.8	5.5	6.9	5.6	5	7.3	5	4.1	4.9	4.7	4.2	4.1	3.9	3.4	
BES 0.7	0.6	0.6	0.6	0.5	0.4	0.6	0.5	0.5	0.4	0.5	0.4	0.4	0.4	0.4	0.4
KAT 0.6	0.5	0.5	0.5	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.5	0.4	0.5	0.4	0.4
BAS 11.5	9.8	9.4	10.6	8.9	7.9	10.9	7.5	7	7.8	7.9	6.8	6.5	6.4	5.7	

Table 2. Computed total annual depositions of lead to six Baltic Sea sub-basins for period 1990-2004. Units: tonnes/year

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004
GUB 118	95	83	65	40	45	56	29	48	52	59	43	39	41	32	
GUF 62	51	43	32	33	27	30	19	24	23	23	21	20	22	16	
GUR 39	32	27	21	19	16	21	13	15	16	16	12	16	16	12	
BAP 416	352	283	266	198	168	216	144	180	190	182	135	163	149	134	
BES 70	46	44	35	29	23	22	20	30	22	24	16	18	19	20	
KAT 64	44	44	29	24	21	18	19	26	23	28	18	21	19	21	
BAS 770	621	525	447	342	299	363	244	322	327	332	245	277	266	235	

Table 3. Computed total annual depositions of mercury to six Baltic Sea sub-basins for period 1990-2004. Units: tonnes/year

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004
GUB	0.9	0.8	0.8	0.8	0.7	0.8	1	0.7	0.8	0.8	0.9	0.7	0.6	0.7	0.6
GUF	0.4	0.3	0.3	0.3	0.3	0.3	0.4	0.3	0.3	0.3	0.3	0.3	0.2	0.3	0.2
GUR	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.1	0.2	0.2	0.1	0.1	0.1	0.1
BAP	2.7	2.4	2.4	2.7	2.3	2.3	2.9	2.2	1.9	2.1	2.1	1.8	1.8	1.7	1.5
BES	0.5	0.4	0.4	0.5	0.4	0.3	0.4	0.3	0.3	0.3	0.3	0.2	0.2	0.2	0.2
KAT	0.4	0.3	0.4	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.2	0.2	0.2	0.2
BAS	5.2	4.4	4.6	4.8	4.2	4.3	5.3	4.1	3.6	3.8	4	3.4	3.3	3.2	2.9

1.1.4 Metadata

1.1.4.1 Technical information:

1. Source:

EMEP/MSC-E

2. Description of data:

The atmospheric depositions of heavy metals were obtained using the latest version of MSCE-HM model developed at EMEP/MSC-E (Travnikov and Ilyin, 2005). The latest available official emission data for the HELCOM countries have been used in the model computations. Emissions of all three metals for each year of this period were officially reported to the UN ECE Secretariat by most of HELCOM countries. These data are available from EMEP emission database WEBDAB: <http://webdab.emep.int>. Some of the countries, in particular, Russia, did not submit the data for 2001. Missing information was obtained on the basis of official data by interpolation. These data were used for modelling of transboundary air pollution by heavy metals within European region including the Baltic Sea and its catchment area.

3. Geographical coverage:

Atmospheric depositions of lead, cadmium, and mercury were obtained for the European region.

4. Temporal coverage:

Timeseries of annual atmospheric depositions are available for the period 1990 – 2004.

5. Methodology and frequency of data collection:

Atmospheric input and source allocation budgets of heavy metals (cadmium, lead, and mercury) to the Baltic Sea and its catchment area were computed using the latest version of MSCE-HM model. MSCE-HM is the regional-scale model operating within the EMEP region. This is a three-dimensional Eulerian model which includes processes of emission, advection, turbulent diffusion, chemical transformations of mercury, wet and dry depositions, and inflow of pollutant into the model domain. Horizontal grid of the model is defined using stereographic projection with spatial resolution 50 km at 60° latitude. The description of EMEP horizontal grid system can be found in the internet (<http://www.emep.int/grid/index.html>). Vertical structure of the model consists of 15 non-uniform layers defined in the terrain-following s-coordinates and covers almost the whole troposphere. Detailed description of the model can be found in EMEP reports (Travnikov and Ilyin, 2005) and in the Internet on EMEP web page <http://www.emep.int> under the link to information on Heavy Metals. Meteorological data used in the calculations for 1990-2004 were obtained using MM5 meteorological data preprocessor on the basis of the Re-analysis project data prepared by National Centers for Environmental Predictions together with National Center of the Atmospheric Research (NCEP/NCAR) in the USA (<http://wesley.ncep.noaa.gov/reanalysis.html>).

Calculations of atmospheric transport and depositions of lead, cadmium, and mercury are provided on the regular basis annually two years in arrears on the basis of emission data officially submitted by Parties to CLRTAP Convention.

1.1.4.2 Quality information:

6. Strength and weakness:

Strength: annually updated information on atmospheric input of lead, cadmium, and mercury to the Baltic Sea and its sub-basins.

Weakness: uncertainties in officially submitted data on emissions of heavy metals.

7. Uncertainty:

The MSCE-HM model has been verified in a number of intercomparison campaigns with other regional models [Sofiev et al., 1996; Gusev et al., 2000; Ryaboshapko et al.,

2001,2005] and has been qualified by means of sensitivity and uncertainty studies [Travnikov, 2000]. It was concluded in this publications that the results of heavy metal airborne transport modeling are in satisfactory agreement with the available measurements and discrepancy does not exceed on average a factor of two. The comparison of calculated versus measured data indicates that the model predicts the observed air concentrations of lead and cadmium within the accuracy of 30%. For concentrations in precipitation the difference between calculated and measured values may reach two times. Computed mercury concentrations deviate from measured values within a factor of two.

The MSCE-HM model results were compared with measurements of EMEP monitoring network [Ilyin and Travnikov, 2006]. The model was evaluated through the comparison with available measurements and the results of other HM models during EMEP TFMM meetings held in 2005 in Zagreb and in Moscow. It was concluded that the MSCE-HM model is suitable for the evaluation of the long range transboundary transport and deposition of HMs in Europe.

8. Further work required:

Further work is required on reducing uncertainties in emission data and modeling approaches used in MSCE-HM model.

1.1.5 References

Gusev A., Ilyin I., Petersen G., van Pul A. and Syrakov D. [2000] Long-range transport model intercomparison studies. Model intercomparison study for cadmium. EMEP/ESC-E Report 2/2000, Meteorological Synthesizing Centre – East, Moscow, Russia. (http://www.msceast.org/reps/2_2000.zip)

Ilyin I. and O. Travnikov (2005) Modelling of heavy metals airborne pollution in Europe: Evaluation of the model performance. EMEP/ESC-E Technical Report 8/2005. (http://www.msceast.org/reps/8_2005.zip)

Ilyin I. and O. Travnikov (2006) Heavy metals: transboundary pollution of the environment. EMEP Status Report 2/2006. (http://www.msceast.org/reps/2_2006.zip)

Ryaboshapko A., Ilyin I., Bullock R., Ebinghaus R., Lohman K., Munthe J., Petersen G., Segneur C., Wangberg I. [2001] Intercomparison study of numerical models for long-range atmospheric transport of mercury. Stage I: Comparison of chemical modules for mercury transformations in a cloud/fog environment. EMEP/ESC-E Technical report 2/2001, Meteorological Synthesizing Centre – East, Moscow, Russia. (<http://www.msceast.org/abstract/201.html>)

Ryaboshapko A., Artz R., Bullock R., Christensen J., Cohen M., Draxler R., Ilyin I., Munthe J., Pacyna J., Petersen G., Syrakov D., and Travnikov O. [2005] Intercomparison study of numerical models for long-range atmospheric transport of mercury. Stage III. Comparison of modelling results with long-term observations and comparison of calculated items of regional balances. EMEP/MSC-E Technical Report 1/2005, Meteorological Synthesizing Centre – East, Moscow, Russia. (http://www.msceast.org/reps/1_2005.zip)

Sofiev M., Maslyayev A. and Gusev A. [1996] Heavy metal model intercomparison. Methodology and results for Pb in 1990. EMEP/MSC-E Report 2/1996, Meteorological Synthesizing Centre - East, Moscow, Russia. (http://www.msceast.org/reps/2_1996.zip)

Travnikov O. [2000] Uncertainty analysis of heavy metals long-range transport modelling. EMEP/MSC-E Technical note 9/2000, Meteorological Synthesizing Centre - East, Moscow, Russia. (<http://www.msceast.org/abstract/900.html>)

For reference purposes, please cite this indicator fact sheet as follows:

[Author's name(s)], [Year]. [Indicator Fact Sheet title]. HELCOM Indicator Fact Sheets 2006. Online. [Date Viewed], http://www.helcom.fi/environment2/ifs/en_GB/cover/.

Last updated 6.9.2006

